

**Impact of universal antiretroviral treatment eligibility on rapid treatment initiation among young adolescents with HIV in sub-Saharan Africa.**

\*Olga Tymejczyk, MPH, Institute for Implementation Science in Population Health, City University of New York, New York, NY, USA

\*Ellen Brazier, MA, Institute for Implementation Science in Population Health, City University of New York, New York, NY, USA

Kara Wools-Kaloustian, MD - Indiana University School of Medicine, Indianapolis, IN, USA

Mary-Ann Davies, MBChB, MMed, PhD, FC - Centre for Infectious Disease Epidemiology and Research, School of Public Health and Family Medicine, University of Cape Town, Cape Town, South Africa

Madeline DiLorenzo, MD, Institute for Implementation Science in Population Health, City University of New York, New York, NY, USA, and Boston Medical Center, Boston, MA, USA

Andrew Edmonds, PhD, MSPH - Department of Epidemiology, The University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

Rachel Vreeman, MD, MS - Department of Pediatrics, Indiana University School of Medicine, Indianapolis, IN, USA

Carolyn Bolton, MBBCh, MSc - Centre for Infectious Disease Research in Zambia, Lusaka, Zambia

Christella Twizere, MD - Centre Hospitalo-Universitaire de Kamenge, Bujumbura, Burundi

Nicollate Okoko, MD - Kenya Medical Research Institute (KEMRI), Nairobi, Kenya

Sam Phiri, PhD, MSc - Lighthouse Trust, Lilongwe, Malawi

Gertrude Nakigozi, MBChB, PhD, MPH- Rakai Health Sciences Program, Kalisizo, Uganda

Patricia Lelo, MD - Kalembelembe Pediatric Hospital, Kinshasa, Democratic Republic of the Congo

Per von Groote, PhD, M.A. - Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland

© The Author(s) 2019. Published by Oxford University Press for the Infectious Diseases Society of America. All rights reserved. For permissions, e-mail: journals.permissions@oup.com.

Annette H. Sohn, MD - TREAT Asia, amfAR – The Foundation for AIDS Research, Bangkok, Thailand  
Denis Nash, PhD, MPH – Institute for Implementation Science in Population Health, City University of  
New York, New York, NY, USA; Department of Epidemiology and Biostatistics, School of Public Health,  
City University of New York, New York, NY, USA  
on behalf of the IeDEA consortium

\*co-first author, contributed equally

SUMMARY (40 words or less): Universal antiretroviral treatment (ART) policies led to increases in rapid  
ART initiation among young adolescents, ages 10-14 years, after enrollment in HIV care. Immediate  
increases in rapid ART initiation were observed after policy adoption in some countries.

CORRESPONDING AUTHOR:

Olga Tymejczyk

Institute for Implementation Science in Population Health, City University of New York

55 W 125<sup>th</sup> St

New York, NY 10027

646-364-9625

ABSTRACT WORD COUNT: 200

TEXT WORD COUNT: 3,498

FOOTNOTES: 0

POTENTIAL CONFLICTS OF INTEREST:

Dr Sohn reports grants from ViiV Healthcare outside of submitted work. Other authors declare no conflicts of interest.

FUNDING:

This work was supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under Award Numbers U01AI096299 (IeDEA Central Africa) and U01AI069924 (IeDEA Southern Africa); the National Institute of Allergy and Infectious Diseases, *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, the National Institute on Drug Abuse, the National Cancer Institute, and the National Institute of Mental Health under Award Number U01AI069911 (IeDEA East Africa). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

CORRESPONDING AUTHOR:

Olga Tymejczyk

Institute for Implementation Science in Population Health, City University of New York

55 W 125<sup>th</sup> St

New York, NY 10027

646-364-9625

## ABSTRACT:

**Background:** Young adolescents with perinatally-acquired HIV are at risk for poor care outcomes. We examined whether universal antiretroviral treatment (ART) eligibility policies (Treat All) improved rapid ART initiation following care enrollment among 10-14-year-olds in seven sub-Saharan African countries.

**Methods:** Regression discontinuity analysis and data for 6,912 10-14-year-old patients were used to estimate changes in rapid ART initiation (within 30 days of care enrollment) following adoption of Treat All policies in two groups of countries: Uganda and Zambia (policy adopted in 2013) and Burundi, Democratic Republic of the Congo, Kenya, Malawi, and Rwanda (policy adopted in 2016).

**Results:** There were immediate increases in rapid ART initiation among young adolescents after national adoption of Treat All. Increases were greater in countries adopting the policy in 2016, compared with those adopting it in 2013: 23.4 percentage points (pp) (95%CI: 13.9-32.8) vs. 11.2pp (95%CI: 2.5-19.9). However, the rate of increase in rapid ART initiation among 10-14-year-olds rose appreciably in countries with earlier treatment expansions, from 1.5pp per year before Treat All to 7.7pp afterwards.

**Conclusions:** Universal ART eligibility has increased rapid treatment initiation among young adolescents enrolling in HIV care. Further research should assess their retention in care and viral suppression under Treat All.

**Keywords:** adolescents, Treat All, ART eligibility, ART initiation, sub-Saharan Africa, regression discontinuity

## INTRODUCTION

In 2018, an estimated 599,000 young adolescents, ages 10 to 14 years, were living with HIV, with close to 90% in sub-Saharan Africa [1, 2]. While data on this age group are limited [3, 4], young adolescents living with HIV are presumed to have acquired HIV perinatally [5, 6]. Studies suggest that adolescents aged 10 to 14 may be less likely than younger children and older age groups to be tested for HIV because of slow-progressing disease, HIV-related stigma, parental concerns about disclosing their own status, and HIV testing strategies, including requirements for caregiver consent, that are not tailored towards adolescents [7-13].

Disproportionately high HIV-related mortality rates have been reported for young adolescents, compared with older adolescents who likely acquire HIV later in life [2]. Young adolescents enrolling in HIV care are often underweight and stunted, and they tend to have lower CD4 counts and more advanced disease than both older adolescents [14, 15] and younger children diagnosed earlier following perinatal infection [16]. Research has shown that adolescents are also at greater risk of failing to initiate antiretroviral treatment (ART), particularly if they are ineligible for treatment at the time they enroll in HIV care [17]. The extent to which treatment eligibility guidelines have constrained ART initiation for adolescents is unknown. Before 2015, the World Health Organization (WHO) provided consolidated guidance on HIV treatment for both adolescents, ages 10 to 19 years, and adults that based ART initiation on CD4 count and clinical eligibility criteria [18-21]. Although a few countries in sub-Saharan Africa extended HIV treatment to all adolescents aged 15 or below in 2013 [22, 23], adolescents only became universally eligible for immediate treatment with the WHO's 2015 recommendation to treat all people living with HIV/AIDS (PLHA), irrespective of immunologic or clinical status [18].

Using longitudinal patient data from seven countries participating in the International epidemiology Databases to Evaluate AIDS (IeDEA) research consortium, we assessed changes in rates of rapid ART initiation among 10-to-14-year-old patients newly enrolling in HIV care following national adoption of Treat All policies for young adolescents or Treat All policies for the general population of PLHA.

## METHODS

### *Data sources and management*

Patient data: The IeDEA consortium ([www.iedea.org](http://www.iedea.org)) assembles sociodemographic and clinical data on adult and pediatric patients receiving HIV care across seven regional cohorts [24]. The data represent diverse clinical sites, the majority of which (87%) are public-sector health facilities, including primary (42%), as well as secondary and tertiary level sites (58%) [25]. In this analysis, we used medical records from 10-to-14-year-old patients newly enrolling in HIV care from 2010 to 2018 in seven sub-Saharan African countries in three IeDEA regions. These countries were selected because patient data were available for analysis following the adoption of universal treatment eligibility policies (Central Africa: Burundi, Democratic Republic of the Congo [DRC], and Rwanda; East Africa: Kenya and Uganda; Southern Africa: Malawi and Zambia). Prior to data analysis, each region's data were standardized by regional data managers in accordance with IeDEA data definitions and formatting standards ([iedeades.org](http://iedeades.org)).

For each country, we identified the date when ART eligibility was first extended to all patients aged 10 to 14 years, either as part of a pediatric (<15 years) Treat All policy, or a general Treat All policy (covering patients of all ages). If a country first adopted a pediatric Treat All policy and subsequently adopted a general Treat All policy, we recorded both dates. We have previously described our systematic search for current and historical ART eligibility guidelines based on publicly available policy

documents, published literature, and input from in-country experts [26]. If the exact day of expansion was unknown, it was assumed to have occurred on the first day of the month in which the policy was adopted.

Data were de-identified before sharing and approved for use by local research ethics committees in each of the leDEA regions included in the study.

#### *Inclusion criteria*

**Patients:** Patients had to be between 10 and 14 years of age at the time of enrollment into HIV care, with at least 30 days of possible follow-up between enrollment and database closure. Patients were excluded if they were known to have transferred to an leDEA site from another clinic or were known to be ART-experienced at enrollment.

**Sites:** Sites had to have patient data available for the period between care enrollment and ART initiation (i.e., pre-ART data) for both ART initiators and non-initiators (i.e., those dying or dropping out of care prior to starting treatment). Sites with data only from the period following ART initiation were excluded.

#### *Outcome and exposure*

The outcome of interest was “rapid” ART initiation, and was defined as initiation of treatment within 30 days of enrollment in HIV care, which is consistent with our prior analyses in adults [27]. This definition differs from the 2017 WHO definition of rapid ART initiation, which is ART-initiation occurring within seven days of HIV diagnosis [28]. The exposure was period of enrollment in HIV care, as defined by the relationship to the calendar date of country-level ART eligibility expansion to Treat All.

#### *Other definitions*

ART was defined as treatment with any regimen of at least three antiretroviral drugs, excluding antiretrovirals taken solely for the prevention of mother-to-child transmission. As a measure of HIV disease severity, pre-treatment CD4 count was defined as the CD4 closest to the enrollment date within a 90-day window (before or after), but no later than one week after ART initiation.

### *Study design*

### *Descriptive analyses*

Patient characteristics, including sex, age, availability of pre-treatment CD4 measure, and median CD4, were described for each country where a pediatric or general Treat All policy extended ART eligibility to all children ages 10 to 14 years, and were aggregated by time period of policy change. The proportion of patients initiating ART rapidly in the year before and after Treat All adoption was calculated for each country and for the Central Africa region (i.e., Burundi, DRC, and Rwanda) because of small sample sizes available for the individual countries.

### *Effect of ART eligibility expansion to Treat All on rapid ART initiation*

The effect of enrollment in HIV care under Treat All on the proportion of young adolescents initiating ART rapidly was assessed using a regression discontinuity design. This approach takes advantage of local randomness in a continuous eligibility assignment variable (calendar time of HIV care enrollment), relative to a cut-off threshold (date of country-level adoption of pediatric or general Treat All). In this quasi-experimental condition, as long as there is no evidence that values of the assignment variable are being manipulated, patients enrolling in care directly before and after the cut-off date are considered exchangeable. Accordingly, there should be no systematic differences in measured and unmeasured



characteristics between the groups, other than the higher probability of treatment eligibility among those enrolling after Treat All policy adoption. If these assumptions are met, observed effects can be interpreted causally, as intention-to-treat estimates [29, 30].

To assess whether there were systematic differences between patients enrolling in HIV care on either side of the threshold, as well as non-random enrollment before or after the Treat All adoption date, we used covariate balance tests and plots of the date of enrollment in HIV care. As complete information about each patient's true ART eligibility status at enrollment prior to Treat All adoption was not known (because of missing information on HIV stage, comorbidities, pregnancy, and/or special population status), the study is an intention-to-treat analysis using a "sharp" regression discontinuity design [29, 30].

We examined the association between calendar time of enrollment in HIV care and rapid ART initiation for two groups of countries: those where a pediatric Treat All policy extended ART eligibility to all children 10 to 14 years old in 2013, and those where a general Treat All policy extended ART eligibility to this age group along with adults in 2016. A discontinuity at the date of each country's Treat All policy adoption allowed for different slopes before and after the cutoff, or threshold, date. Local linear regression models [31] were used to estimate predicted outcomes and risk differences at the Treat All threshold date as follows:

$$E[Y_i|Z_i] = \beta_0 + \beta_1 * Z_i + \beta_2 * 1[Z_i \geq 0] + \beta_3 * Z_i * 1[Z_i \geq 0]$$

where  $Y_i$  is the patient-level outcome (rapid ART initiation),  $Z_i$  is the number of days between a patient's enrollment date and national Treat All policy adoption date (negative if patient enrolled before the policy was adopted), and  $1[Z_i \geq 0]$  indicates whether a patient enrolled after the policy was adopted or not.

Data-driven Imbens-Kalyanaraman bandwidths [32] were used to define windows of time around the date of Treat All adoption within which predicted outcomes and risk differences at the Treat All threshold date were estimated. All observations within the bandwidth were weighted equally. Sensitivity analyses were completed using three other bandwidth sizes, ranging from 150 to 450 days.

In the countries with a general Treat All expansion to all ages following a pediatric Treat All policy, an additional regression discontinuity analysis for the general Treat All adoption was completed to assess whether further expansions of eligibility criteria to encompass older age groups affected rapid ART initiation among already-eligible 10-to-14-year-olds. Such effects could be positive (increase in rapid ART initiation because of, for example, stigma reduction) or negative (decrease in rapid ART initiation because of, for example, facility capacity constraints).

#### *Trends in rapid ART initiation before and after Treat All adoption*

To characterize trends in rapid ART initiation following enrollment into HIV care, slopes from linear regression models for the period before and after the date of Treat All adoption were compared, and expressed as percentage point change in rapid ART initiation per year (i.e., average annual rate of increase).

Analyses were completed in SAS 9.4 and Stata/MP version 15.1.

## RESULTS

### *Sample characteristics*

Longitudinal data were available for 7,296 patients ages 10 to 14 years who enrolled in HIV care between 2010 and 2018, including 7,239 (99.2%) with at least 30 days of possible follow-up. Of these patients, 6,912 (95.5%) had no evidence of transfer from another site or ART prior to enrollment.

Among the seven countries in the analysis, five (Burundi, DRC, Kenya, Malawi, Rwanda) adopted general Treat All policies that extended treatment eligibility to young adolescents ages 10 to 14 years in 2016. Two countries (Uganda, Zambia) adopted pediatric Treat All policies in 2013, which extended treatment to all children younger than 15 years, and adopted general Treat All policies in 2016. (Supplementary Table 1)

Among the 6,912 patients who met study inclusion criteria, 3,592 (52.0%) were in countries where pediatric Treat All was adopted in 2013, and 3,320 (48.0%) in countries where general Treat All was adopted in 2016. The median age at enrollment in HIV care was 12 years (interquartile range [IQR]: 11-13), with no significant age differences before and after Treat All adoption (Supplementary Table 2), and 58.3% were female. The availability of pre-treatment CD4 count measures varied across countries and before vs. after Treat All policy adoption. Few patients had pre-treatment CD4 measures following the adoption of Treat All policies, particularly in countries that introduced the policy in 2016 (20.1% overall, and only 3% among patients from Malawi). Among patients with a pre-treatment CD4 count, the median value before Treat All adoption was 315 cells/ $\mu$ l (IQR: 124-551) in countries adopting general Treat All policies in 2016 and 363 cells/ $\mu$ l (IQR: 193-589) in countries that adopted a pediatric Treat All policy in 2013 (Table 1).

**Table 1. Characteristics of adolescents 10 to 14 years of age enrolling in HIV care (n=6,912), 2010-2018.**

	General Treat All adopted in 2016						Pediatric Treat All adopted in 2013		
	Burundi	DRC	Kenya	Malawi	Rwanda	Overall	Uganda	Zambia	Overall
Treat All adoption date	Sep 2016	Sep 2016	Jul 2016	May 2016	Jul 2016	2016	Dec 2013	Dec 2013	2013
Total enrollments (%)	178 (2.6%)	123 (1.8%)	1,752 (25.3%)	1,168 (16.9%)	99 (1.4%)	3,320 (48.0%)	461 (6.7%)	3,131 (45.3%)	3,592 (52.0%)
Period of enrollment									
Before Treat All adoption	154 (86.5%)	110 (89.4%)	1548 (88.4%)	1067 (91.4%)	82 (82.8%)	2,961 (89.2%)	257 (55.7%)	1635 (52.2%)	1,892 (52.7%)
After Treat All adoption	24 (13.5%)	13 (10.6%)	204 (11.6%)	101 (8.6%)	17 (17.2%)	359 (10.8%)	204 (44.3%)	1496 (47.8%)	1,700 (47.3%)
Sex									
Male	73 (41.0%)	67 (54.5%)	702 (40.1%)	511 (43.8%)	50 (50.5%)	1,403 (42.3%)	186 (40.3%)	1294 (41.3%)	1,480 (41.2%)
Female	105 (59.0%)	56 (45.5%)	1050 (59.9%)	657 (56.3%)	49 (49.5%)	1,917 (57.7%)	275 (59.7%)	1837 (58.7%)	2,112 (58.8%)
Age (years)									
Median (IQR)	12 (11-13)	12 (11-13)	12 (11-13)	12 (11-13)	12 (11-13)	12 (11-13)	12 (11-13)	12 (11-13)	12 (11-13)
Pre-treatment CD4 measure before Treat All adoption									
CD4 measure available	42 (27.3%)	79 (71.8%)	1,073 (69.3%)	235 (22.0%)	66 (80.5%)	1,495 (50.5%)	167 (65.0%)	1,090 (66.7%)	1,257 (66.4%)
Median CD4 count (cells/ $\mu$ l) (IQR)	417 (220-762)	297 (93-499)	315 (120-574)	308 (145-456)	400 (176-675)	315 (124-551)	377 (197-671)	359 (193-579)	363 (193-589)
Pre-treatment CD4 measure after Treat All adoption									
CD4 measure available	5 (20.8%)	1 (7.7%)	53 (26.0%)	3 (3.0%)	10 (58.8%)	72 (20.1%)	106 (52.0%)	704 (47.1%)	1,700 (47.7%)
Median CD4 count (cells/ $\mu$ l) (IQR)	N/A	N/A	397 (233-558)	N/A	N/A	N/A	408 (194-579)	343 (173-545)	347 (173-554)

IQR = interquartile range; DRC = Democratic Republic of the Congo

Accepted Manuscript

Distributions of baseline characteristics among newly enrolling patients were similar just before and just after Treat All adoption (Supplementary Table 2). No major discontinuity in the number of new enrollments around the date of Treat All adoption was observed (Supplementary Figure 1).

**Figure 1. Proportions of adolescents 10 to 14 years old initiating ART rapidly (within 30 days of enrollment in HIV care) in the years before and after Treat All adoption, by country or region.**

*Rapid ART initiation before and after Treat All adoption (descriptive analysis)*

There were appreciable increases in rapid ART initiation among young adolescents in the year following Treat All adoption in all seven countries. Increases ranged from 16.3 percentage points (pp) in Zambia (from 37.4% in the year before to 53.7% in the year after) to 33.7pp in Uganda (from 41.3% to 75.0%, respectively). The proportion of young adolescents rapidly initiating ART in the year after Treat All adoption was highest in Malawi and Kenya (88.1% and 86.5%, respectively), both of which adopted a general Treat All policy in 2016. (Figure 1).

*Effect of Treat All adoption on rapid ART initiation (regression discontinuity analysis)*

Statistically significant increases in rapid ART initiation among young adolescents were observed immediately after national Treat All policies expanded treatment eligibility for this age group. A larger absolute effect was observed in the group of countries that adopted a general Treat All policy in 2016 (Burundi, DRC, Kenya, Malawi, Rwanda): 23.4pp (95% CI: 13.9, 32.8), compared with the countries with a 2013 pediatric Treat All policy (Uganda, Zambia), where there was an 11.2pp increase in rapid ART

initiation (95% CI: 2.5, 19.9). In the 2016 Treat All group, 85.4% of young adolescents enrolling immediately after Treat All adoption initiated ART rapidly (up from 62.0% immediately before), while in the 2013 group, 50.2% of young adolescents initiated rapidly after the expansion (up from 39.0% immediately before; Table 2, Figure 2).

**Table 2. Effect of ART eligibility expansion to treat all adolescents 10 to 14 years old and trends in rapid ART initiation before and after Treat All adoption.**

Expansion date	2016	2013	2016
Expansion type	General Treat All	Pediatric Treat All	General Treat All
Countries	Burundi, DRC, Kenya, Malawi, Rwanda	Uganda, Zambia	Uganda, Zambia
<hr/>			
Risk difference at the Treat All adoption threshold*	23.4	11.2	-1.1
95% CI	13.9 - 32.8	2.5 - 19.9	-13.9 - 11.7
p-value	<0.001	0.012	0.862
Imbens-Kalyanaraman bandwidth, days	681	780	252
N within bandwidth	970	1,937	665
Predicted outcomes at the Treat All threshold*			
Enrollment just before Treat All adoption	62.0%	39.0%	77.7%
Enrollment just after Treat All adoption	85.4%	50.2%	76.6%
Relative change after Treat All adoption	37.7%	28.7%	-1.4%
Slopes before and after Treat All adoption**			
Percentage point change in rapid ART initiation per year before Treat All adoption	4.1	1.5	5.8

Percentage point change in rapid ART initiation			
per year after Treat All adoption	2.0	7.7	4.4
P-value for difference of slopes	0.693	<0.001	0.928

---

ART = antiretroviral treatment; DRC = Democratic Republic of the Congo

\*Risk difference and predicted outcomes at the Treat All threshold are from regression discontinuity analyses.

Effects are calculated at the guideline expansion threshold of one day before versus one day after Treat All adoption.

\*\*Slope comparison is from separate linear regression models comparing the periods before Treat All adoption and after Treat All adoption.

**Figure 2. Trends in rapid ART initiation before and after Treat All adoption among adolescents 10 to 14 years old, by year of Treat All adoption.**

There was no statistically significant change in rapid ART initiation among young adolescents immediately after general Treat All policies were adopted in Uganda and Zambia in 2016 (Table 2). Pediatric Treat All policies were already in place in these two countries, and the proportion of young adolescents initiating ART rapidly was 77.7% immediately before the general Treat All policy, versus 76.6% immediately after.

Results of sensitivity analyses using other bandwidths were consistent with the findings based on the data-driven Imbens-Kalyanaraman bandwidth (Supplementary Table 3).

*Trends in rapid ART initiation before and after Treat All adoption (slope comparison)*

The average annual rate of increase in rapid ART initiation among young adolescents rose in countries where a pediatric Treat All policy was adopted in 2013, from 1.5pp per year before Treat All adoption to



7.7pp afterwards (Table 2). However, no statistically significant change in the annual rate of increase in rapid ART initiation was observed in countries that adopted a general Treat All policy in 2016. In addition, no rate change was observed in Uganda and Zambia following the expansion of pediatric Treat All policies to include all PLHA in 2016.

## DISCUSSION

Whether part of a pediatric Treat All policy or a general Treat All policy, expansions of HIV treatment eligibility to those under age 15 were followed by significant and substantial increases in ART initiation among 10- to 14-year-olds within 30 days of enrollment in HIV care. Increases in rapid ART initiation were particularly large following national adoptions of a general Treat All policy (i.e., for all ages) in 2016 in Burundi, DRC, Kenya, Malawi, and Rwanda. Observed increases in the proportion of young adolescents rapidly initiating ART following national adoption of Treat All policies may have substantial clinical importance, given evidence indicating that young adolescents with perinatally-acquired HIV often enroll in care late and do not initiate ART until they are at advanced stages of disease [14, 15, 33-35].

Pre-treatment CD4 measures were not available for a large proportion of patients, making it difficult to ascertain clinical eligibility for treatment among those enrolling before Treat All policies were adopted. However, among the 56.7% of young adolescents with pre-treatment CD4 data available prior to Treat All adoption, median CD4 counts for each country in our study were well below 500 cells/ul, the previous immunologic threshold for treatment eligibility. This suggests that there were gaps in rapid ART initiation for young adolescents when CD4-based eligibility criteria were in effect. These findings are in accordance with previous research that has highlighted a range of social and structural barriers to HIV

care and treatment for adolescents, whose needs may not be adequately met by service delivery strategies designed for younger children or adult HIV patients. [6, 7, 36, 37].

The absolute effect of treatment eligibility expansions for young adolescents was greater under general Treat All policies than under earlier pediatric Treat All policies. This was noteworthy, given that large numbers of adult patients, newly-eligible under Treat All, could potentially strain HIV service provision, leading to the crowding out of vulnerable groups, such as young adolescents, for whom tailored interventions and services are recommended [11, 15, 38-40]. These results lend support for the supposition that general Treat All policies, with harmonized treatment recommendations for different population groups, are easier to implement in real-world treatment settings than prior policies targeting specific groups. Age-agnostic guidelines may help simplify the provision of HIV treatment in low-resourced health systems in ways that lead to efficiency gains in service delivery [41, 42].

Given the three-year interval between the pediatric Treat All expansions of 2013 and the general Treat All expansions of 2016, findings may also reflect temporal trends in provider preparedness, health system capacities to rapidly implement expanded treatment guidelines, and improved strategies for initiating patients on ART, despite decreases in donor funding for HIV during the period [43, 44]. While immediate increases in rapid ART initiation among 10- to 14-year-olds were smaller following national adoptions of pediatric Treat All policies, the average annual rate of change in rapid ART initiation increased significantly following the adoption of these policies in 2013. Consistent with the smaller immediate effect observed at the Treat All adoption threshold, this may reflect a gradual roll-out or delayed implementation of the policy. In contrast, the lack of a statistically significant change in annual rates of rapid ART initiation following general Treat All policy adoptions in 2016 may be due to the high rates of rapid ART initiation achieved immediately after the policy (85.4%), with limited space for further increases above this level, as well as regression to the mean.

A concerning finding was the large decrease in pre-treatment CD4 measurement, as reflected by an increased proportion of young adolescents with no CD4 measures prior to treatment initiation. This finding is not unique to our study population [26, 45, 46], and reflects a larger trend of abandoning the use of CD4 monitoring altogether in sub-Saharan Africa, driven by combinations of cost and laboratory supply chain issues, as well as prioritization of viral load over CD4 testing [47-49]. Treat All policies have negated the need for CD4 to assess eligibility for treatment, and routine CD4 monitoring after ART initiation is generally not necessary in virally suppressed patients in settings with routine viral load testing [50, 51]. However, assessing *pre-treatment* CD4 remains important for identifying severely immunodeficient individuals who need enhanced clinical services, such as treatment of opportunistic infections [52-54] and for monitoring progress towards achieving the public health goals of HIV care and treatment scale-up [51].

A strength of this analysis is the use of a regression discontinuity design with real-world service delivery data from diverse settings in seven sub-Saharan African countries that adopted Treat All policies at two points in time. This quasi-experimental design provides support for the causal interpretation of the association between expanded ART eligibility under Treat All and increases in rapid ART uptake among young adolescents newly enrolling into HIV care. The use of a data-driven Imbens-Kalyanaraman bandwidth [32] and sensitivity analyses with three other bandwidths enabled us to generate robust effect estimates with minimal risk of researcher bias.

A limitation of our study is lack of complete data on treatment eligibility criteria for patients enrolling prior to national Treat All policy adoption (e.g., pre-treatment CD4 counts, WHO staging, and coinfection with TB). Such data would have allowed for the use of a “fuzzy” regression discontinuity design [29] to better reflect the probabilistic distribution of ART eligibility in the pre-Treat All sample. In addition, the

limited availability of data on patient characteristics beyond age, sex, and pre-treatment CD4 count restricted our ability to assess whether patients on each side of the regression discontinuity threshold were similar with respect to other pre-treatment covariates. We also lacked data on service delivery strategies, including tailored services for adolescents, and supply side constraints, such as drug stockouts, that may influence rapid ART initiation among young adolescents enrolling into care. Moreover, while we know the dates when Treat All policies were adopted in each country, lags in site-level implementation likely varied across sites and countries included in this analysis [55]. Finally, the use of a 30-day rapid ART initiation window, intended to enable comparisons with this group's prior work [27], limits the comparability of findings to WHO's rapid ART initiation estimates defined by a 7-day window after confirming HIV diagnosis.

These limitations notwithstanding, our results suggest that expanded treatment eligibility under Treat All has benefited 10-to-14-year-olds by getting them onto treatment more rapidly following enrollment in care. As there are few age-disaggregated data related to the HIV care continuum for this age group, this study fills an important gap, indicating that an increasing share of young adolescents may be initiating ART rapidly under Treat All policies. While these results are encouraging, further research is needed on effective strategies for enrolling children with perinatally-acquired HIV in HIV care earlier and improving care retention and ART adherence among adolescent patients to support sustained viral suppression among this vulnerable population.

#### ACKNOWLEDGMENTS

The authors thank patients, providers, and administrative staff at participating leDEA facilities. We also are grateful for information on ART eligibility expansions contributed by Dr Edith Apondi, Dr Lastone

Chitembo, Dr Nathan Ford, Dr John Humphrey, Dr Olivia Keiser, Yee Yee Kuhn, Malango Msukwa, Dr Martina Penazzato, and Dr Ellon Twinomuhwezi.

## REFERENCES

1. United Nations Children's Fund (UNICEF). Children, HIV and AIDS: The world in 2030. New York, NY, **2018**.
2. Slogrove AL, Sohn AH. The global epidemiology of adolescents living with HIV: time for more granular data to improve adolescent health outcomes. *Current Opinion in HIV and AIDS* **2018**; 13:170-8.
3. Slogrove AL, Mahy M, Armstrong A, Davies M-A. Living and dying to be counted: What we know about the epidemiology of the global adolescent HIV epidemic. *Journal of the International AIDS Society* **2017**; 20:4-15.
4. Idele P, Gillespie A, Porth T, et al. Epidemiology of HIV and AIDS Among Adolescents: Current Status, Inequities, and Data Gaps. *JAIDS Journal of Acquired Immune Deficiency Syndromes* **2014**; 66:S144-S53.
5. Ng'eno BN, Kellogg TA, Kim AA, et al. Modes of HIV transmission among adolescents and young adults aged 10-24 years in Kenya. *International journal of STD & AIDS* **2018**; 29:800-5.
6. Lowenthal ED, Bakeera-Kitaka S, Marukutira T, Chapman J, Goldrath K, Ferrand RA. Perinatally acquired HIV infection in adolescents from sub-Saharan Africa: a review of emerging challenges. *The Lancet Infectious diseases* **2014**; 14:627-39.
7. Enane LA, Davies M-A, Leroy V, et al. Traversing the cascade: urgent research priorities for implementing the 'treat all' strategy for children and adolescents living with HIV in sub-Saharan Africa. *Journal of virus eradication* **2018**; 4:40-6.
8. Davies M-A, Kalk E. Provider-Initiated HIV Testing and Counselling for Children. *PLoS medicine* **2014**; 11:e1001650.

9. Chikwari CD, Dringus S, Ferrand RA. Barriers to, and emerging strategies for, HIV testing among adolescents in sub-Saharan Africa. *Current Opinion in HIV and AIDS* **2018**; 13:257-64.
10. Govindasamy D, Ferrand RA, Wilmore SM, et al. Uptake and yield of HIV testing and counselling among children and adolescents in sub-Saharan Africa: a systematic review. *Journal of the International AIDS Society* **2015**; 18:20182.
11. Lightfoot M, Dunbar M, Weiser SD. Reducing undiagnosed HIV infection among adolescents in sub-Saharan Africa: Provider-initiated and opt-out testing are not enough. *PLoS medicine* **2017**; 14:e1002361.
12. Zandoni BC, Elliott RJ, Neilan AM, Haberer JE. Screening for HIV and linkage to care in adolescents: insights from a systematic review of recent interventions in high- versus low- and middle-income settings. *Adolescent health, medicine and therapeutics* **2018**; 9:211-35.
13. Wong VJ, Murray KR, Phelps BR, Vermund SH, McCarragher DR. Adolescents, young people, and the 90–90–90 goals: a call to improve HIV testing and linkage to treatment. *Aids* **2017**; 31:S191-S4.
14. Koech E, Teasdale CA, Wang C, et al. Characteristics and outcomes of HIV-infected youth and young adolescents enrolled in HIV care in Kenya. *AIDS (London, England)* **2014**; 28:2729-38.
15. Apondi E, Humphrey JM, Sang E, et al. Trends Over Time for Adolescents Enrolling in HIV Care in Kenya, Tanzania, and Uganda From 2001–2014. *JAIDS Journal of Acquired Immune Deficiency Syndromes* **2018**; 79:164-72.
16. Westerlund E, Jerene D, Mulissa Z, Hallström I, Lindtjørn B. Pre-ART retention in care and prevalence of tuberculosis among HIV-infected children at a district hospital in southern Ethiopia. *BMC Pediatrics* **2014**; 14:250.
17. Desmond S, Tanser F, Vreeman R, et al. Access to antiretroviral therapy in HIV-infected children aged 0–19 years in the International Epidemiology Databases to Evaluate AIDS

(IeDEA) Global Cohort Consortium, 2004–2015: A prospective cohort study. *PLoS medicine* **2018**; 15:e1002565.

18. World Health Organization (WHO). Guideline on when to start antiretroviral therapy and on pre-exposure prophylaxis for HIV. Geneva: World Health Organization, **2015**.

19. World Health Organization (WHO). Consolidated guidelines on the use of antiretroviral drugs for treating and preventing HIV infection: Recommendations for a public health approach - Second edition. Geneva: WHO, **2016**.

20. World Health Organization (WHO). Antiretroviral therapy for HIV infection in adults and adolescents: recommendations for a public health approach. (2010 revision). Geneva: World Health Organization, **2010**.

21. World Health Organization (WHO). Consolidated guidelines on the use of antiretroviral drugs for treating and preventing HIV infection: recommendations for a public health approach. Geneva: World Health Organization, **2013**.

22. Ministry of Health (Republic of Uganda). Addendum to the Antiretroviral Treatment Guidelines for Uganda, **2013**.

23. Ministry of Health (Government of the Republic of Zambia). Zambia Consolidated Guidelines for the Prevention and Treatment of HIV Infection, **2013**.

24. IeDEA. International Epidemiologic Databases to Evaluate AIDS. Available at: <https://www.iedea.org/>. Accessed June 1, 2018.

25. Fritz CQ, Blevins M, Lindegren ML, et al. Comprehensiveness of HIV care provided at global HIV treatment sites in the IeDEA consortium: 2009 and 2014. *Journal of the International AIDS Society* **2017**; 20.

26. Tymejczyk O, Brazier E, Yiannoutsos C, et al. HIV treatment eligibility expansion and timely antiretroviral treatment initiation following enrollment in HIV care: A metaregression analysis of programmatic data from 22 countries. *PLoS medicine* **2018**; 15:e1002534.


27. Tymejczyk O, Brazier E, Yiannoutsos CT, et al. Changes in rapid HIV treatment initiation after national "treat all" policy adoption in 6 sub-Saharan African countries: Regression discontinuity analysis. *PLoS medicine* **2019**; 16:e1002822.
28. World Health Organization (WHO). Guidelines for managing advanced HIV disease and rapid initiation of antiretroviral therapy, July 2017. Geneva: World Health Organization, **2017**.
29. Moscoe E, Bor J, Barnighausen T. Regression discontinuity designs are underutilized in medicine, epidemiology, and public health: a review of current and best practice. *Journal of clinical epidemiology* **2015**; 68:122-33.
30. Bor J, Fox MP, Rosen S, et al. Treatment eligibility and retention in clinical HIV care: A regression discontinuity study in South Africa. *PLoS medicine* **2017**; 14:e1002463.
31. Hahn J, Todd P, Van der Klaauw W. Identification and estimation of treatment effects with a regression-discontinuity design. *Econometrica* **2001**; 69:201-9.
32. Imbens G, Kalyanaraman K. Optimal bandwidth choice for the regression discontinuity estimator. *The Review of economic studies* **2012**; 79:933-59.
33. Kariminia A, Law M, Davies M-A, et al. Mortality and losses to follow-up among adolescents living with HIV in the IeDEA global cohort collaboration. *Journal of the International AIDS Society* **2018**; 21:e25215-e.
34. Koller M, Patel K, Chi BH, et al. Immunodeficiency in children starting antiretroviral therapy in low-, middle-, and high-income countries. *Journal of acquired immune deficiency syndromes (1999)* **2015**; 68:62-72.
35. The Collaborative Initiative for Paediatric HIV Education Research (CIPHER), Slogrove AL, Schomaker M, et al. The epidemiology of adolescents living with perinatally acquired HIV: A cross-region global cohort analysis. *PLoS medicine* **2018**; 15:e1002514.
36. World Health Organization (WHO). HIV and adolescents: guidance for HIV testing and counselling and care for adolescents living with HIV: recommendations for a public health approach and considerations for policy-makers and managers, **2013**.



37. Mark D, Armstrong A, Andrade C, et al. HIV treatment and care services for adolescents: a situational analysis of 218 facilities in 23 sub-Saharan African countries. *Journal of the International AIDS Society* **2017**; 20:21591-.
38. Kose J, Tiam A, Ochuka B, et al. Impact of a Comprehensive Adolescent-Focused Case Finding Intervention on Uptake of HIV Testing and Linkage to Care Among Adolescents in Western Kenya. *Journal of acquired immune deficiency syndromes (1999)* **2018**; 79:367-74.
39. Simms V, Dauya E, Dakshina S, et al. Community burden of undiagnosed HIV infection among adolescents in Zimbabwe following primary healthcare-based provider-initiated HIV testing and counselling: A cross-sectional survey. *PLoS medicine* **2017**; 14:e1002360.
40. MacPherson P, Munthali C, Ferguson J, et al. Service delivery interventions to improve adolescents' linkage, retention and adherence to antiretroviral therapy and HIV care. *Tropical medicine & international health : TM & IH* **2015**; 20:1015-32.
41. Hontelez JA, Chang AY, Ogbuaji O, de Vlas SJ, Barnighausen T, Atun R. Changing HIV treatment eligibility under health system constraints in sub-Saharan Africa: investment needs, population health gains, and cost-effectiveness. *Aids* **2016**; 30:2341-50.
42. Ford N, Penazzato M, Vitoria M, et al. The contribution of observational studies in supporting the WHO 'treat all' recommendation for HIV/AIDS. *J Virus Erad* **2018**; 4:5-8.
43. Kates J, Wexler A, Leif E. Donor government funding for HIV in low- and middle-income countries in 2016: . Menlo Park, CA: Kaiser Family Foundation & UNAIDS, **2017**.
44. Kates J, Wexler A, Lief E. Financing the response to HIV in low- and middle-Income countries: International assistance from donor governments in 2015. Menlo Park, CA: Kaiser Family Foundation and UNAIDS, **2016**.
45. The IeDEA and COHERE Cohort Collaborations. Global Trends in CD4 Cell Count at the Start of Antiretroviral Therapy: Collaborative Study of Treatment Programs. *Clinical Infectious Diseases* **2018**; 66:893-903.


46. Teasdale CA, Yuengling K, Preko P, et al. Persons living with HIV with advanced HIV disease: need for novel care models. *Journal of the International AIDS Society* **2018**; 21:e25210-e.
47. Ford N, Meintjes G, Vitoria M, Greene G, Chiller T. The evolving role of CD4 cell counts in HIV care. *Current Opinion in HIV and AIDS* **2017**; 12:123-8.
48. Tenforde MW, Walker AS, Gibb DM, Manabe YC. Rapid antiretroviral therapy initiation in low- and middle-income countries: A resource-based approach. *PLoS medicine* **2019**; 16:e1002723-e.
49. Ford N, Meintjes G, Pozniak A, et al. The future role of CD4 cell count for monitoring antiretroviral therapy. *Lancet Infect Dis* **2015**; 15:241-7.
50. Ford N, Meintjes G, Pozniak A, et al. The future role of CD4 cell count for monitoring antiretroviral therapy. *The Lancet infectious diseases* **2015**; 15:241-7.
51. Nash D, Robertson M. How to Evolve the Response to the Global HIV Epidemic With New Metrics and Targets Based on Pre-Treatment CD4 Counts. *Current HIV/AIDS reports* **2019**; 16:304-13.
52. Hakim J, Musiime V, Szubert AJ, et al. Enhanced Prophylaxis plus Antiretroviral Therapy for Advanced HIV Infection in Africa. *N Engl J Med* **2017**; 377:233-45.
53. World Health Organization. Guidelines for managing advanced HIV disease and rapid initiation of antiretroviral therapy. , **2017**.
54. Phillips A, Shroufi A, Vojnov L, et al. Sustainable HIV treatment in Africa through viral-load-informed differentiated care. *Nature* **2015**; 528:S68-76.
55. Brazier E, Maruri F, Duda SN, et al. Implementation of “Treat-all” at adult HIV care and treatment sites in the Global IeDEA Consortium: results from the Site Assessment Survey. *Journal of the International AIDS Society* **2019**; 22:e25331.

Figure 1



Accepted Manuscript

Figure 2



Accepted